



(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:
21.10.1998 Bulletin 1998/43

(51) Int. Cl.⁶: A61K 47/48

(21) Application number: 98105208.7

(22) Date of filing: 02.05.1986

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(30) Priority: 03.05.1985 US 730682

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
92201393.3 / 0 506 207
86903757.2 / 0 221 176

(71) Applicant:
Access Pharmaceuticals, Inc.
Dallas, Texas 75207 (US)

(72) Inventor: Allen, Larry M.
Denver, Colorado 80206 (US)

(74) Representative:
Allard, Susan Joyce et al
BOULT WADE TENNANT,
27 Fumival Street
London EC4A 1PQ (GB)

Remarks:

This application was filed on 23 - 03 - 1998 as a
divisional application to the application mentioned
under INID code 62.

(54) Pharmaceutical vehicles for reducing transdermal flux

(57) The present invention provides a topical pharmaceutical composition comprising a pharmacologically active agent and a vehicle characterized in that said composition includes a potentiator for the pharmacologically active agent which comprises a water soluble zinc-containing compound that provides zinc ions that can complex or chelate with the pharmacologically active agent.

Description

Field of the Invention

- 5 Method of inducing reservoir effect in skin and mucous membranes so as to increase penetration and residence time of pharmacologically active agents therein.

Background of the Invention

- 10 There are many localized disease conditions which are effectively treated by topical application of suitable physiological agents. In order for such treatments to be maximally effective, it is necessary that as much of the pharmacologically active agent as possible be absorbed into the skin where it can make contact with the disease condition in the dermal tissues without being lost by rubbing off on clothing or evaporation. At the same time, the agent must not penetrate so effectively through the skin as to be rapidly lost to the lymphatic and vascular circulatory systems. This latter factor is especially important when the pharmacologically active agent is toxic when used systemically.

The ideal vehicle for topically applied pharmaceuticals is therefore one which can produce a "reservoir effect" in the skin or mucous membranes to which the topical treatment is applied. This "reservoir effect" is defined as an enhancement of the skin or membrane's ability to both absorb and retain pharmacologically active agents, i.e., to increase skin or membrane residence time, decrease drug transit time and reduce transdermal flux.

- 20 A number of compounds are known to enhance the ability of pharmacologically active agents to penetrate the skin and mucous membranes, for example, N-bis-azacyclopentan-2-onyl-alkanes, 1-substituted azacycloheptan-2-ones and higher alkyl-substituted azacyclopentan-2-ones, as well as dimethylsulfoxide and lower alkyl sulfoxides. These compounds, however, have the disadvantage of allowing rapid systemic dispersion of the pharmacologically active agents away from the localized site of pathology. Many topical medicaments, such as the retinoids used in the treatment of acne, and methotrexate, used in the treatment of psoriasis, are systemically toxic. The retinoids, for example, are known to cause damage to unborn fetuses. Thus, there is a need for a method of enhancing the ability of such medicaments to penetrate into the skin or mucous membrane so that a lesser total dosage may be used, while at the same time retarding their ability to move from the skin to the interior of the body.

- 30 The problem posed by the paradoxical requirement that a systemically toxic topical medicament be efficiently absorbed into, but not through, the skin, has heretofore gone unrecognized. There have been no vehicle additives available to the pharmacological industry which act to enhance the "reservoir capacity" of the skin and mucous membranes so that the amount of pharmacologically active agents reaching and being retained at the site of localized pathologies is maximized.

35 Summary of the Invention

This invention is a method for inducing a reservoir effect in skin and mucous membranes so as to enhance penetration and retention of topically applied pharmacologically active therapeutic and cosmetic agents therein. The invention also relates to topical treatment methods involving such reservoir effect enhancers, and to pharmaceutical compositions containing them.

- 40 The additives of this invention are water-soluble zinc-containing compounds, preferably zinc halide, zinc sulfate, zinc nitrate, zinc acetate, and/or zinc stearate, and most preferably zinc chloride.

- The pharmacologically active agents with which the water-soluble zinc-containing compounds are used are preferably those containing hydroxyl, oxo, sulphydryl, amine, carboxyl, and other anionic groups in configurations which readily allow complexation or chelation with zinc ions.

- 45 In inducing a reservoir effect in the skin and mucous membranes to reduce transdermal flux so that drugs are absorbed and retained therein in larger amounts for longer periods of time than has heretofore been possible, the water-soluble zinc-containing compounds of this invention act as potentiators for the pharmacologically active agents. Potentiation is defined as overcoming or reducing undesirable effects such as systemic toxicity and extending the range of effectiveness of the pharmacologically active agent, or both.

Description of the Drawing

- 50 The drawing is a graph of test results showing the skin reservoir effect achieved by the method of this invention utilizing nordihydroguaiaretic acid with zinc chloride, as compared to the same compound without zinc chloride. The nordihydroguaiaretic acid was labelled with carbon-14 for radiotracer analysis. The graph shows immediate absorption and longer retention of larger amounts of the nordihydroguaiaretic acid with zinc chloride present than without. The availability of nordihydroguaiaretic acid, a lipoxigenase inhibitor, to therapeutically act upon localized pathologies, both with

and without added zinc chloride, is measured by the areas under the respective curves. It is apparent that skin bioavailability is greatly enhanced in the presence of a zinc-containing compound. The drug flux rate can be calculated as a function of the area under the curve by dividing this area by dose. As is apparent, drug flux rate is substantially decreased over the entire dosage range by the addition of zinc chloride.

Detailed Description of the Invention

The zinc-containing compounds of this invention are generally any water-soluble organic or inorganic zinc salts which dissociate in the topical vehicle so as to provide zinc ions for complexation or chelation with the pharmacologically active agents present in the vehicle. Examples of suitable zinc-containing compounds are zinc halide, zinc sulfate, zinc nitrate, zinc acetate, and/or zinc stearate. The most preferred zinc-containing compound is zinc chloride.

Such water-soluble zinc-containing compounds may be prepared by means known to the art, and many are commercially available.

The topical preparations of this invention having enhanced reservoir inducing capacities may be prepared by mixing a water-soluble zinc-containing compound into the pharmaceutical preparation or vehicle in which a reservoir-inducing capacity is to be created by means known to the art so as to promote complexation or chelation of the pharmacologically active agents therein with zinc ions. The presence of other metallic ions which would unfavorably compete with zinc for complexation or chelation sites is to be avoided. The zinc salt can also be mixed with the pharmacologically active agent to form a chelate or complex which is then incorporated into the pharmaceutical carrier.

The zinc-containing compounds are preferably present in an equimolar ratio with the pharmacologically active agents, so as to cause maximum complexation or chelation. Where stratum corneum destruction, i.e. decorrification, is desirable, an excess of such zinc-containing compounds which also act as escharotics, e.g. zinc chloride, may be used. (Generally, concentrations of 35% (0.257 moles per 100 grams) or more zinc chloride will cause tissue destruction when topically applied.) Normally, use of equimolar concentrations of zinc chloride and the pharmacologically active agent will not involve the use of escharotic amounts of zinc chloride, however less than 35% zinc chloride should be considered an upper limit when no escharotic effect is desired. Less than an equimolar ratio of zinc-containing compound to pharmacologically active agent may be employed where it is not desired that all the medicament be absorbed into the skin or mucous membranes, e.g. in connection with mouthwash and douche preparations where attack on free-swimming organisms is also desirable.

Other ingredients may be added to the preparations, including coloring agents, stability-enhancing agents, antioxidants, and the like. Preferably these additives will not compete with the pharmacologically active agents for zinc; however when necessary, excess zinc-containing compounds may be used to compensate for the zinc complexing or chelating effect of such additives.

The pharmacologically active agents of this invention are those intended for topical application to achieve localized therapeutic or cosmetic effects. A partial list of suitable pharmacologically active agents includes steroids, antifungals, anti-unicellular microorganism agents, antiviral agents, antiparasitic agents, antineoplastic agents, anti-leprosy agents, antimetabolites, cell-regulatory agents, immuno-pharmacological agents, allergens, antihistaminic agents, anti-inflammatory agents, anesthetic agents, analgesic agents, counter-irritants, vitamins, nutrients, diagnostic agents, radiopaque agents, cryoprotective agents, perfumes, insect repellants, hair dyes, anticarsing agents, sun screens, melanin-stimulating agents, antiperspirants, antisecretory agents, depilatories, hair restorers, wrinkle-reducing agents, antidandruff agents, emollients, rubifacients, and cosmetic agents in general.

The mechanism by which the reservoir-inducing effect of this invention is produced is not known; however, it is preferred that the pharmacologically active agents contain hydroxy, oxo, sulphydryl, amine, carboxyl, or other anionic groups, or combinations thereof, in conformations which allow complexation and/or chelation by zinc ions.

Preferred pharmacologically active agents of this invention are:

1. Antineoplastic agents including NDGA (nordihydroguaiaretic acid), VP-16 (epipodophyllotoxin beta-D ethylidene glucopyranoside- ϵ -toposide), VM-26 (epipodophyllotoxin beta-D thenylidene glucopyranoside- ϵ -toposide), 4-demethyl epipodophyllotoxin, diethylstilbestrol, dihydranthranol, cyclophosphamide, mitomycin, daunomycin, platinum cis-diamine-dichloride, adriamycin, allolpurinol, 5-fluorouracil, and methotrexate.

2. Immunopharmacological agents which may be topically applied including polypeptide nanoparticles comprising interleukin or active fragments thereof, antibodies or active fragments thereof, interferons, and liposomes. Such delivery systems providing sustained release of pharmacologically active agents are effectively localized or held in place by zinc according to this invention.

3. Steroids, which are utilized for a wide range of therapeutic purposes including anti-inflammation, antipruritic, enhancement of moisture retention, etc., including: dexamethasone, hydrocortisone, hydrocortisone acetate, hydroxy hydrocortisone, hydrocortisone valerate, triamcinolone acetonide, triamcinolone hexacetonide, amcinonide, fluocinolone acetonide, fluocinonide, flurandrenolide, diflurasone diacetate, betamethasone dipropionate,

betamethasone, betamethasone benzoate, betamethasone valerate, halcinonide, desoximethasone, desonide, prednisolone, and clocortolone pivalate.

4. Antifungal agents which are used to treat fungus infections on the skin, hair, and nails, such as athlete's foot (*tinea pedis*), jock itch (*tinea cruris*), ringworm (*tinea corporis*), which can be caused by a number of fungi, particularly *Trichophyton rubrum*, *Trichophyton mentagrophytes*, and *Epidermophyton floccosum*, and *Microsporum canis*. These antifungal agents include haloprogin, itochloro, miconazole nitrate, tolnaftate, thiabendazole, chloroxine, amphotericin, candicidin, fungimycin, nystatin, chlordanol, clotrimazole, ethonam nitrate, miconazole nitrate, pyrrol-nitrin, lezatione, ticlatone, tolnaftate, triacetin, carbonic acid derivatives; dithiocarbamate, thiourea, thiocyanates; aromatic carboxylic acids and the amides thereof, benzoic acid, salicylic acid, salicylic acid amide and anilide; aromatic sulfides, polysulfides, and sulfoxides, 5,5-dichloro-2,2-dihydroxydiphenylsulfide; invert soaps, quaternary ammonia and phosphonium compounds, decamethylene-bis-(4-thio-pyridine-methyl-tosylate); quinoline derivatives, 8-hydroxyquinoline sulfate, halogenated quinolines, 7-iodo-8-hydroxy-quinoline-5-sulfonic acid, 5-chloro-7-iodo-8-hydroxy-quinoline, 5-chloro-8-hydroxy-quinoline, 5,7-dichloro-8-hydroxyquinoline, 5,7-diiodo-8-hydroxy-quinoline, decamethylene-bis (4-amino-quinolium chloride); benzothiazole derivatives, (2-dimethylamino-6-(beta-diaminoethoxy)-benzothiazole dihydrochloride); imidazole derivatives, 1-(o-chloro-alpha-alpha-diphenyl-benzyl)-imidazole, 1-[o,p-dichloro-beta-(o,p-dichlorobenzoyloxy)-phenethylimidazole]; benzimidazole derivatives, 2-phenyl-benzimidazole, 2-furfurylbenzimidazole; thiazine derivatives, 3,5-dibenzyltetrahydro-1,3,5-thiadiazine-2-thione; furan derivatives, 5-nitro-2-furfuryl-3-chloropropionate; quinones, tetrachloro-p-benzoquinone, 1,4-naphthoquinone, phenanthraquinone; sulfonamides and sulfones; aromatic diamidines, 2-hydroxysilbamidine, and diamidinodiphenylamine.

Antifungal agents are also used to treat vaginal infections caused by *Candida albicans* and related yeasts; these agents include dioctyl sodium sulfosuccinate, haloprogin, miconazole nitrate, potassium sorbate, propionate salts, such as calcium propionate and sodium propionate, and sodium lauryl sulfate.

5. Antibacterial agents, which are utilized for treating skin infections such as impetigo, ecthyma, folliculitis, boils, and acute prurynchia, and for treating skin wounds and as a wound cleanser, and which may be used in this invention, including sulfonamides, penicillins, cephalosporins, penicillinase, lincomycins, vancomycins, tetracyclines, chloramphenicol, and streptomycins; including within this group the following compounds: gramicidin, neomycin, polymyxin beta sulfate, tetracycline, benzethonium chloride, gentamicin sulfate, nitrofurazone, benzalkonium chloride, hexylresorcinol, chloroxylenol, cloflucaban, carboxylic acid (phenol), triclocaban, and triclosan.

6. Antiviral agents, including those used to treat warts, such as glacial acetic acid, ascorbic acid, calcium pantothenate, lactic acid, salicylic acid, cantharidin, and podophyllin; and antiviral agents used to treat cold sores or herpes simplex such as acyclovir, benzalkonium chloride, alcohol, allantoin, anhydrous glycerin, benzocaine, camphor, carbamide peroxide, lanolin, menthol, petrolatum, and phenol; and antiviral agents including those used to treat herpes genitalis such as urea, idoxuridine, amantadine, methisazone, cytarabine, interferons, chloroform, ether, bacillus calmette-guerin, and levamisole.

7. Antiparasitic agents including anthelmintic agents (agents that destroy or expel intestinal worms) capable of penetrating the skin of the animal to be treated, e.g. benzimidazole compounds, tetramisole, levamisole, and isoquinoline compounds.

8. Pediculicides, for mites (or scabies) and lice, including lidane, pyrethrins, piperonyl butoxide, malathion, and crotamiton.

9. Acne treatment compounds including benzoyl peroxide, resorcinol, resorcinol monoacetate, sulfur, povidone-iodine, salicylic acid, phenol, fluocinolone acetonide, para-aminobenzoic acid, sodium thiosulfate, meclocycline sulfosalicylate, sodium sulfacetamide, tetracycline hydrochloride, aliphatic dicarboxylic acids, e.g. adipic and azelaic acids, and sulfurated lime.

10. Antipsoriasis agents including cytostatic agents, which retard skin-cell growth; keratolytic agents, which loosen and dissolve scales, tar preparations, whose mode of action is uncertain; hydrocortisone preparations, which reduce itching and inflammation; anti-itch preparations; and antimicrobials. These antipsoriasis agents include coal tar preparations, juniper tar, pine tar, allantoin, saponated creol, menthol, mercury oleate, phenol preparations, resorcinol, salicylic acid, anthralin, and methotrexate.

11. Leprosy agents including 4,4'-diaminodiphenyl sulfone.

12. Anesthetic agents for pain and itching, inflamed skin, sunburn, insect bites, burns, wounds, hemorrhoids, poison ivy, poison oak, including: benzocaine, lidocaine, lidocaine hydrochloride, dibucaine, dibucaine hydrochloride, procaine, tetracaine, tetracaine hydrochloride, tronothane, dydonine, dydonine hydrochloride, pramoxine hydrochloride, benzyl alcohol, diperoon, butamben picrate, cyclomethacaine sulfate, and dimethisoquin hydrochloride.

13. Analgesic agents for pain and itching, inflamed skin, sunburn, insect bites, burns, wounds, hemorrhoids, poison ivy, poison oak, including: salicylic acid derivatives; N,N-dimethyl aspartic acid; N,N-dimethyl glutamic acid, triamine salicylate, methyl salicylate; antipyrine, aspirin, and salicylamide.

14. Counter-irritants (agents applied locally to produce an inflammatory reaction with the object of distracting and

relieving a deep seated inflammatory process) including methyl salicylate, camphor, menthol, eugenol, eucalyptol, thymol, allyl isothiocyanate (mustard oil), capsicum preparations, histamine dihydrochloride, methyl nicotinate, and turpentine oil.

15. Antihistamines which are used principally against itching, but are also mildly anesthetic, including diphenhydramine hydrochloride, phenyltoloxamine dihydrogen citrate, pyrilamine maleate, triptelenamine hydrochloride.

16. Diagnostic agents including allergenic extracts for diagnosis and immunotherapy of specific allergy offenders from the following categories: pollens, foods, dusts, epidermals, insects and stinging insects, fungi, molds, yeasts; tests for sensitivity to therapeutic penicillin (benzyl-penicillioyl-polylysine); tests for sensitivity to tetanus antigens, diphtheria antigens, streptococcus antigens, tuberculin, Candida antigens, Trichophyton antigens, and Proteus antigens.

17. Vitamins and nutrients for skin, hair, and scalp conditions, including anti-scarring agents, vitamins B₃, B₆, A, D, and E.

18. Cosmetic agents and perfumes including compositions to reduce the appearance of wrinkles such as water soluble elastin and pregnenolone; skin depigmenting agents and bleaches, including hydroquinone and monobenzone.

19. Sunscreens including: dioxybenzone, oxybenzone, padimate O, padimate A, aminobenzoic acid, cinoxate, diethanolamine p-methoxycinnamate, ethyl 4-[bis(hydroxypropyl)] aminobenzoate, ethylhexyl salicylate, glyceryl aminobenzoate, homosalate, the combination of lawsone and dihydroxyacetone, red petrolatum, and sulisobenzonzone.

The foregoing are simply examples of pharmacologically active agents including therapeutic and cosmetic agents which may be used with enhanced effectiveness for their known properties in accordance with this invention.

In addition to the reservoir effect produced by zinc ions in skin and mucous membranes, a direct potentiating effect has been observed when pharmaceutical preparations containing zinc ions and pharmacologically active agents are injected directly into diseased tissues, particularly solid tumors. The mechanism for this potentiating effect is not known; however, it may be caused by a reservoir-inducing effect directly on the tissues involved.

Dosage forms for topical application may include lotions, ointments, creams, gels, suppositories, nasal solutions, mouthwashes, sprays, aerosols and the like. Typical carriers which make up the foregoing dosage forms include water, acetone, isopropyl alcohol, stearyl alcohol, freons, ethyl alcohol, polyvinyl pyrrolidone, propylene glycol, polyethylene glycol, fragrances, gel-producing materials, mineral oil, stearic acid, spermaceti, sorbitan, monooleate, polysorbates, "Tweens," sorbitol, methyl cellulose, etc.

The amount of the composition, and thus of the pharmacologically active agent therein to be administered, will obviously be an effective amount for the desired result expected therefrom. This, of course, will be ascertained by the practitioner utilizing his ordinary skill. Due to enhanced activity which is achieved, the dosage of agent may often be decreased from that generally applicable. In accordance with usual prudent formulating practices, a dosage near the lower useful range of the particular agent may be employed initially and the dosage increased as indicated from the observed response.

Preferred compositions are illustrated in the following examples:

40 EXAMPLE 1

Antineoplastic Preparation

Two compositions utilizing zinc chloride, nordihydroguaiaretic acid (NDGA), acid (EDTA), butylated hydroxytoluene (BHT), stearyl alcohol, purified water, polyethylene glycol having an average molecular weight of 400 (PEG 400), and polyethylene glycol having an average molecular weight of 3350 (PEG 3350) were prepared in the following manner: the purified water was placed in a clean glass container of suitable capacity; the water was heated to about 80-90°C with stirring; and zinc chloride was added to the heated water, continuing the stirring until the zinc chloride dissolved. The ethylenediaminetetraacetic acid was next slowly added with mixing until dissolved. In a separate glass container of suitable size, the polyethylene glycol 400 was heated to about 80-90°C with stirring; the NDGA was added thereto; then the BHT; and this mixture was added to the zinc chloride/ethylenediaminetetraacetic acid solution with stirring. The entire mixture was then cooled to about room temperature and passed through a number 3 roller mill until smooth. The polyethylene glycol 3350 was then heated to about 80-90°C in a suitable container and the milled ingredients added thereto with mixing.

The final compositions in w/wt % were as follows:

TABLE 1

Composition	Compound	
	A	B
zinc chloride	29.8	10.0
NDGA	4.6	4.6
EDTA	14.7	4.93
BHT	1.1	1.1
stearyl alcohol	0.5	0.5
H ₂ O	18.3	18.3
PEGO 400	26.4	26.4
PEGO 3350	4.5	4.5

EXAMPLE 2

Reservoir Effect

This study was designed to provide basic pharmacokinetic data on the disposition of Carbon 14 (¹⁴C) labelled nor-dihydroguaiaretic acid (NDGA) applied dermally in modified compounds A and B described in Example 1. In addition, the distribution of zinc was measured for the dermally applied Compound A.

The ¹⁴C-NDGA compound exhibited a specific radioactivity of 20.2 Ci/mole (66.9 micro Ci/mg) and a purity of 96.9% by mass spectrometry and by radioautography of thin-layer chromatography plates developed in benzene:isopropanol:acetic acid:water (25:5:2:10).

Subsequently, 25.1 mg of the ¹⁴C-NDGA-Compound (66.9 micro Ci/mg) were mixed with 12.35 g of Compound A. Analyses of triplicate samples of the final mixture for ¹⁴C by counting and for NDGA by high-pressure liquid chromatography (HPLC) demonstrated homogeneity, with a content of 51 micro g of ¹⁴C-NDGA compound/mg of Compound A. The specific radioactivity of the NDGA was 3.00×10^3 micro Ci/micro g.

Similarly, 26.3 mg of the original ¹⁴C-NDGA compound (66.9 micro Ci/mg) were mixed with 12.55 g of Compound A devoid of Z and EDTA to obtain a mixture for the study of the dermal penetration of NDGA from Compound A devoid of Z and EA. Analyses of triplicate samples of the modified Compound A showed the final mixture to be homogeneous with regard to ¹⁴C and NDGA; it contained 53 micro g of ¹⁴C-NDGA compound/mg of vehicle. The specific radioactivity of the diluted NDGA was 3.41×10^{-3} micro Ci/micro g.

The compounds were dermally applied to young adult Sprague-Dawley rats by the following protocol: under ether anesthesia, the back skin of the rat was prepared by removing the hair from a 5 x 5-cm area with a clipper and the residual hair stubble was removed with a wax depilatory. Then the skin was stripped repeatedly (5x) with adhesive tape until the stratum corneum was removed. Then 0.5 gm of the formulation was weighed on a 5 x 5-cm sheet of polypropylene, which was applied to the prepared skin. It was secured in place by hypoallergenic tape. Finally, the bandage was over-wrapped with bandage tape. After treatment, the rats were caged individually in metabolism cages, which allowed free access to food and water and provided for separate collection of urine and feces.

The testing of Compound A with ¹⁴C-NDGA was performed in 15 male Sprague-Dawley rats (mean weight 339 ± 16 g). They received an average of 0.520 (± 0.032) g of Compound A containing ¹⁴C-NDGA. The mean dose of ¹⁴C-NDGA was 78.5 (± 7.0) mg/kg of body weight. The rats were housed individually in metabolism cages providing for free access to food and water and for the separate collection of urine and feces. Groups of 3 rats were sacrificed at 4, 24, 48, 72, and 96 hr and excreta were collected from each rat during 24-hr periods. In addition to the usual collection of tissues, the skin site of application was excised after wiping the site with water-moistened tissue. The wipes were added to the wrappings, which were immersed in a small container of acetone.

The testing of Compound A devoid of Zn and EDTA was performed on 15 male Sprague-Dawley rats (mean weight 241 ± 7 g). They received an average of 0.390 (± 0.019) g of Zn-free C205 containing ¹⁴C-NDGA. The average dose of ¹⁴C-NDGA was 83.2 mg/kg of body weight. Groups of three rats were bled terminally and tissues were taken at 4, 24, 48, 72, and 96 hr after dosing. At each sacrifice time, those three rats scheduled to be sacrificed next were also bled nonterminally from the orbital sinus. The wrappings and wipes of the skin site were taken at the time of sacrifice and added to acetone as described above.

EP 0 872 248 A2

The results of the study are given in Table 2A for Compound A containing ^{14}C -NDGA and Table 2B for Zn-Free Compound A containing ^{14}C -NDGA. The results of analysis for tissue distribution of zinc as a percent of dose in rats receiving Compound A containing ^{14}C -NDGA are given in Table 2C.

TABLE 2A

Tissue Distribution of ^{14}C as a Percent of Dose in Rats Receiving Compound A Containing ^{14}C -NDGA, Dermal					
Tissue	Mean (\pm S.D.) ^a Percent of the Dose of ^{14}C Found at Hours				
	4	24	48	72	96
Organs (%)	4.62 (± 2.27)	7.55 (± 1.85)	10.29 (± 8.8)	10.43 (± 6.94)	13.45 (± 3.45)
Skinsite (%)	20.8 (± 8.4)	19.6 (± 5.2)	11.1 (± 1.4)	8.58 (± 5.41)	7.01 (± 2.28)
^a N = 3					

TABLE 2B

Tissue Distribution of ^{14}C as a Percent of Dose in Rats Receiving Zn-Free Compound A Containing ^{14}C -NDGA, Dermal					
Tissue	Mean (\pm S.D.) ^a Percent of the Dose of ^{14}C Found at Hours				
	4	24	48	72	96
Organs (%)	10.28 (± 6.12)	11.08 (± 9.65)	10.47 (± 10.33)	5.41 (± 4.21)	8.03 (± 1.48)
Skinsite (%)	4.15 (± 0.76)	12.20 (± 5.9)	7.86 (± 3.75)	4.72 (± 3.18)	1.43 (± 0.55)
^a N = 3					

TABLE 2C

Tissue Distribution of Zn as a Percent of Dose in Rats Receiving Compound A, Dermal					
Tissue	Mean Percent of the Dose of Zn found at Hours				
	4	24	48	72	96
Organs (%)	3.28	6.54	6.79	11.47	12.69
Skinsite (%)	10.5	11.8	9.58	6.46	4.99
^a N = 3					

The study was continued for testing Compound B with ^{14}C -NDGA; Zn- and EDTA-free Compound B; and modified Compound B with no BHT and 0.10 EDTA. The following Table 2D lists the compositions of the compounds and the amounts of materials used for preparing the compounds containing ^{14}C -NDGA. These compounds were analyzed for ^{14}C by scintillation counting and for NDGA by liquid chromatography.

TABLE 2D

Constituent	Composition (%)		
	Compound B Formula- tion	Zn-Free Compound B Formulation	Modified Compound B Formulation
Compound Zn	10.00	0.00	10.00
Compound EDTA	4.93	0.00	0.10
Compound NDGA	4.60	4.60	4.60
Compound BHT	1.10	1.10	0.00
Water, purified	18.32	18.32	19.42
PEG 400	2.60	14.50	14.19
PEG 8000	53.45	49.39	46.69
Stearyl Alcohol	5.00	12.09	5.00
	100%	100%	100%
Compounds containing ¹⁴ C-NDGA:			
mg of ¹⁴ C-NDGA	25.75	25.40	25.20
g of Formulation	12.55	12.55	12.75
% NDGA in final mixture	4.80	4.79	4.78

The mean rat body weights, average doses of the formulations, and mean doses of ¹⁴C-NDGA in mg/kg of body weight for the three current protocols were: 297 ± 15 g (standard deviation), 512 ± 28 mg, and 82.7 ± 2.0 mg/kg for Compound B; 325 ± 12 g, 570 ± 26 mg, and 84.0 ± 1.4 mg/kg for Zn-free Compound B; and 328 ± 27 g, 575 ± 45 mg, and 84.2 ± 2.8 mg/kg for modified Compound B. Fifteen rats were used for each study and groups of three rats were sacrificed at 4, 24, 48, 72 and 96 hr after dosing. At each time, blood, liver, skin site, intestines and contents, carcass, and combined wrappings and wipes were collected. Also, from the groups sacrificed at 24, 48, 72, and 96 hr urine, feces, and cage washings were collected.

The results of the study are given in Table 2E for Compound B; Table 2F for Zn-free Compound B; and Table 2G for modified Compound B.

TABLE 2E

Tissue Distribution of ¹⁴ C as a Percent of Dose in Rats Receiving Compound B Containing ¹⁴ C-NDGA, Dermal					
Tissue	Mean (± S.D.) ^a Percent of the Dose of ¹⁴ C Found at Hours				
	4	24	48	72	96
Organs (%)	1.92 (±1.34)	4.41 (±1.44)	4.31 (±2.5)	3.38 (±0.52)	1.87 (±1.1)
Skinsite (%)	16.3 (±9.2)	11.9 (±6.1)	8.92 (±4.10)	6.55 (±1.76)	5.11 (±3.01)
^a N = 3					

TABLE 2F

Tissue Distribution of ^{14}C as a Percent of Dose in Rats Receiving Zn-Free Compound B Containing ^{14}C -NDGA, Dermally					
Tissue	Mean (\pm S.D.) ^a Percent of the Dose of ^{14}C Found at Hours				
	4	24	48	72	96
Organs (%)	1.16 (± 0.92)	2.29 (± 2.26)	2.38 (± 1.73)	7.39 (± 10.05)	7.73 (± 10.9)
Skinsite (%)	2.21 (± 1.44)	4.45 (± 4.36)	6.07 (± 2.90)	18.6 (± 6.3)	12.0 (± 4.99)
^a N = 3					

TABLE 2G

Tissue Distribution of ^{14}C as a Percent of Dose in Rats Receiving Modified Compound B Containing ^{14}C -NDGA, Dermally					
Tissue	Mean (\pm S.D.) ^a Percent of the Dose of ^{14}C Found at Hours				
	4	24	48	72	96
Organs (%)	4.35 (± 3.06)	1.65 (± 0.68)	4.41 (± 3.57)	2.54 (± 2.05)	0.97 (± 0.19)
Skinsite (%)	14.4 (± 7.1)	23.0 (± 5.5)	17.1 (± 4.7)	16.3 (± 0.6)	17.7 (± 8.2)
^a N = 3					

The above results show that the addition of the water-soluble zinc-containing compound causes the organic molecule to more quickly absorb into the skin in larger quantities, and to be retained in the skin longer than when the zinc-containing compound is not present.

EXAMPLE 3

Antineoplastic Enhancement

For the organic compounds listed, in the case of those having two or more benzene rings, .0145 moles per 100 cc's was used, and for those having one or no benzene rings, .029 moles per 100 cc's was used. The organic compound was measured into a clean vial and PEGO 400 was added with mixing until dissolved.

For the test compounds containing zinc chloride, zinc chloride was first dissolved in the PEGO 400 to prepare a stock solution containing 0.69% zinc chloride, and this solution was added with mixing to the vials containing the organic compounds being tested.

These compounds were tested with and without zinc chloride for their effectiveness as antitumor agents against xenografts of human breast adenocarcinoma, MX-1, grown in athymic (nude) mice of Balb/c background by intratumor injection according to the following protocol: each animal was inoculated intradermally on the dorsum near the nape of the neck with 0.05 ml of an MX-1 tumor homogenate. Tumor weights, in milligrams, were calculated from a measurement of the length, width and height in millimeters of the tumors using the formula $(L \times W \times H)/2$. The animals were randomized in groups to ensure representation of smaller and larger tumors. The tumors were treated by intratumor injection with 0.05 ml of each test composition. Each composition was tested utilizing five animals. The animals were treated only once. Results are set forth in Table 3.

TABLE 3

	Organic Compound Known Anti-Cancer Agent	Tumor Free 60 Days	Premature Death	Tumor at Death	Tumor Recur- rence
5	VP-16 (no Zn)	2	2	3	0
	VP-16 + Zn	5	0	0	0
10	*1/5 VP-16 + Zn	3	0	2	2
	VM-26 (no Zn)	0	0	5	0
	VM-26 + Zn	5	0	0	0
	*1/5 VM-26 + Zn	5	0	0	0
15	4'-demethylepipodophyllotoxin (no Zn)	1	0	4	3
	4'-demethylepipodophyllotoxin + Zn	5	0	0	0
	diethylstilbestrol (no Zn)	0	2	5	0
20	diethylstilbestrol + Zn	3	0	2	1
	dithranol (no Zn)	1	0	4	0
	dithranol + Zn	4	0	1	1
	cyclophosphamide (no Zn)	0	0	5	0
25	cyclophosphamide + Zn	3	2	0	0
	mitomycin (no Zn)	1	4	2	0
	mitomycin + Zn	5	0	0	0
30	daunomycin (no Zn)	3	2	5	3
	daunomycin + Zn	5	0	0	0
	platinum cis-diaminedichloride (no Zn)	1	0	4	0
	platinum cis-diaminedichloride + Zn	5	0	0	0
35	adriamycin (no Zn)	0	1	2	4
	*1/10 adriamycin (no Zn)	2	1	2	2
	adriamycin + Zn	4	0	1	1
40	*1/10 adriamycin + Zn	1	3	1	1
	allopurinol (no Zn)**	0	--	5	--
	*5/2 allopurinol (no Zn)	0	0	5	0
	allopurinol + Zn	1	0	4	4
45	*5/2 allopurinol + Zn	1	0	4	4

* Dosage level decreased or increased as indicated

** All sacrificed on day 25.

These results show the reduced toxicity and enhanced antineoplastic effectiveness achieved by the addition of zinc ions.

EXAMPLE 4**Antineoplastic Potentiation**

Compositions and organic compounds not previously known as antineoplastic agents were prepared with and with-

out zinc chloride according to the procedure of Example 3 and tested for their ability to eradicate tumors following the protocol described in Example 3. Results are set forth in Table 4.

TABLE 4

Organic Compound	Tumor Free 60 Days	Premature Death	Tumor at Death	Tumor Recurrence
3-tertbutylphenol (no Zn)	1	3	1	0
3-tertbutylphenol + Zn	5	0	0	0
4-tertbutylphenol (no Zn)	5	0	0	0
4-tertbutylphenol + Zn	5	0	0	0
p-hydroxycinnamic acid (no Zn)	1	0	4	0
p-hydroxycinnamic acid + Zn	4	1	0	0
norisoguaiacin (no Zn)	2	1	1	0
norisoguaiacin + Zn	5	0	0	0
di-NDGA (no Zn)	4	0	1	0
di-NDGA + Zn	5	0	0	0
azelaic acid (no Zn)	1	0	4	0
azelaic acid + Zn	5	0	0	0
1-(3,4-diacetoxyphenyl)-4-phenylbuta-1,3-diene (no Zn)	1	0	4	0
1-(3,4-diacetoxyphenyl)-4-phenylbuta-1,3-diene + Zn	3	0	2	2
1,4-bis(3,4-dihydroxyphenethyl)benzene (no Zn)	2	0	3	3
1,4-bis(3,4-dihydroxyphenethyl)benzene + Zn	2	0	3	3

These results show that the antineoplastic activity of drugs can be increased to worthwhile levels while at the same time reducing toxicity utilizing zinc ions.

EXAMPLES 5-10

Enhancement of Topical Antineoplastic Agents

In Examples 5-10, wherein reference is made to the testing of mixtures for antitumor activity against B-16 melanoma and Sarcoma-180 solid tumor growth in mice, the following procedures were utilized. To the extent that a particular example modified the procedure, such modification will be indicated in the particular example.

Both types of tumors were grown intradermally or subcutaneously in the mice. The B-16 melanoma was grown in BDF₁ mice and the S-180 tumor was grown in ICR mice. Each mouse was injected intradermally with about 0.01 ml of a saline suspension containing about 1×10^6 cells of the tumor cells per 0.01 ml into a preshaven area on the back of the neck of the mouse. The tumors were allowed to grow until they had an approximate size of about 25-100 mg, calculated by the length of the tumor multiplied by the width and height of the tumor measured in millimeters and dividing the product by two. On the first day of treatment, the animals with tumor sizes outside of the size range were culled and the remaining animals were randomly divided into control and test groups. When the tumors had reached the appropriate size, usually at about day six, the tumors were punctured uniformly and then treated with either a test compound or a control by topical application to the surface of the tumor. Generally, two topical applications were made 24 hours apart. The materials were applied to obtain from about a 1 to about 2 mm coating over the surface of the tumor. The animals were thereafter observed and their weights and the size of their tumors were periodically measured.

The results of each of the experiments include the following:

- (a) the starting number (n) of animals within a treatment group of an experiment;
 (b) the average tumor size in milligrams of the animals treated with the mixture and the average tumor size of the control animals;
 (c) the ratio multiplied by 100 of the average size of the tumors of the treated animals to that of the control animals (T/C), wherein T = average size of treated mice and C = average tumor size of control mice;
 (d) the percentage of both treated and control animals clear of tumor; and
 (e) the percentage of animals of the original number surviving.

The later three measurements for a particular experiment were all taken at the same time and range generally from 21 to about 33 days after tumor inoculation. A T/C value of 42 or less is indicative of activity. In all of the following tables for Examples 5-10, the control results are given in parenthesis ().

EXAMPLE 5

Two formulations of meso-NDGA were prepared in a PEGO base. Their compositions are as set forth in Table 5.

TABLE 5

Mixture	NDGA (meso)	H ₂ O	EtOH	PEGO 3350
1	3.6	24.3	48.5	23.4
2	6.9	0	0	93.1

Mixture No. 1 was prepared by dissolving the NDGA in absolute ethanol by warming and stirring; thereafter, the water was added slowly to the NDGA solution. The mixture was heated to evaporate sufficient solvent to obtain a mixture of about 130% of the weight of the NDGA, and was then incorporated into the PEGO base. Mixture No. 2 was made by simply dissolving the NDGA in the PEGO base with warming and stirring.

EXAMPLE 6

The NDGA formulations of Example 5 were tested for potential antitumor activity against B-16 melanoma grown in mice. The procedure utilized was that previously described. The results are given below in Table 6.

TABLE 6

n	T/C ¹	Tumor Size (Control)	% Clear (Control)	% Survival (Control)
9	87	954±698 (1091±547)	0 (0)	33 (80)
9	114	748±621 (645±335)	0 (0)	77 (77)
10*	24	138±99 575±270	20 (0)	100 (60)

¹T/C ratio was calculated between days 21-24.

*Animal treated once via a 0.05 ml. intratumor injection on day 6 after tumor inoculation

EXAMPLE 7

Several formulations of zinc chloride in a PEGO base were prepared by first dissolving the zinc chloride in water and then mixing the zinc chloride solution into the PEGO base. The formulations had approximately the following weight/weight percent compositions as set forth in Table 7.

TABLE 7

Mixture	ZnCl ₂	H ₂ O	PEGO
3	41.9	11.6	46.5

EP 0 872 248 A2

TABLE 7 (continued)

Mixture	ZnCl ₂	H ₂ O	PEGO
4	30	16	53.8
5	14.6	4.1	81.3
6	28.6	7.9	63.5
7	46.1	13.8	40
8	15	8	77
9	13.8	9.2	77
10	5.5	3.7	90.8

EXAMPLE 8

Mixtures of Example 7 were tested for potential antitumor activity against B-16 melanoma and S-180 solid tumor grown in mice in accordance with the procedures previously described. The results are given in Table 8.

TABLE 8

S-180					
Mixture	n	T/C ¹	Tumor Size (Control)	% Clear (Control)	% Survival (Control)
3	8	51	559±476 (1095±360)	25 (10)	100 (100)
3	10	40	366±345 (934±656)	11 (10)	90 (100)
6	8	48	524±462 (1095±360)	12 (10)	100 (100)
6	10	115	1074±687 (934±656)	10 (10)	100 (100)
7	5	0	0 (752±511)	100 (0)	80 (100)
7	10	0	0 (550±184)	100 (0)	90 (100)
7	10	0	0 (997±421)	100 (0)	100
10	8	74	815±472 (1095±360)	0 (10)	87 (100)
B-16					
Mixture	n	T/C ¹	Tumor Size (Control)	% Clear (Control)	% Survival (Control)
3	10	6	162±253 (2505±1844)	60 (0)	100 (100)
4	10	16	106±114 (644±342)	30 (0)	100 (70)
5	10	47	1191±764 (2505±1844)	22 (0)	90 (100)
6	10	4	94±217 (2502±1844)	80 (0)	100 (100)
7	9	15	169±216 (1091±547)	37 (0)	88 (80)
7	9	0	0 (654±335)	100 (0)	88 (77)
7	10	2	31±100 (1805±968)	90 (0)	100 (80)
8	10	43	277±209 (644±342)	10 (0)	40 (70)
9	10	1.0	25±63 (1733±2254)	80 (70)	90 (80)

¹T/C ratio calculated between days 20-23.

²T/C ratio calculated between days 21 and 25.

EXAMPLE 9

Mixtures of zinc chloride, EDTA and NDGA were prepared and formulated in a PEGO base. The mixtures were pre-

pared by dissolving the NDGA and EDTA in a portion of the PEGO base by warming and stirring until dissolved. The zinc chloride was dissolved in water and warmed. The warm zinc chloride solution was added to the warm PEGO containing the NDGA and EDTA and stirred until cooled to room temperature. The composition of the mixtures is given in approximate weight/weight percentage as set forth in Table 9.

TABLE 9

Mixture	ZnCl ₂	NDGA	EDTA	H ₂ O	PEGO
11	27.5	6.9	14.7	18.3	32.6
12	28	6.8 ¹	14.7	18.2	32.9

¹ - T/C ratio calculated at day 24 except for Mixture 11 which was calculated at day 21.

EXAMPLE 10

The mixtures of Example 9 were tested for their potential antitumor activities against B-16 melanomas grown in mice in accordance with the procedure previously described. The results are given in Table 10.

TABLE 10

B-16					
Mixture	n	T/C ¹	Tumor Size (Control)	% Clear (Control)	% Survival (Control)
11	10	8	51±118 (711±286)	70 (0)	100 (100)
12	10	0	0 (711±286)	60 (0)	100 (100)

¹ - T/C ratio calculated at day 24 except for Mixture 11 which was calculated at day 21.

The foregoing Examples 5-10 show the potentiating effect of zinc chloride on antineoplastic agents topically applied, showing improvement over the antineoplastic activity of zinc chloride alone, and comparable amounts of NDGA even when the NDGA is injected into the tumor.

EXAMPLE 11

Appropriate human and animal models are chosen for the disease conditions treated by the following compounds as hereinabove described, and the compounds tested to determine effective and toxic dosages with and without the addition of equimolar amounts of zinc chloride, zinc iodide, zinc bromide, zinc sulfate, zinc nitrate, zinc stearate, and zinc acetate. From the results the Therapeutic Index, equivalent to ratio of toxic to effective dosage is calculated, and this Index for the compounds with and without zinc additives compared to show an increase in Therapeutic Index for compounds containing zinc additives as compared to compounds without such additives: NDGA (nordihydroguaiaretic acid), VP-16 (epipodophyllotoxin beta-D ethylidene glucopyranoside-etoside), VM-26 (epipodophyllotoxin beta-D thenylidene glucopyranoside-tenoside), 4'-demethyl epipodophyllotoxin, diethylstilbestrol, dihydran, cyclophosphamide, mitomycin, daunomycin, platinum cis-diamine-dichloride, adriamycin, allopurinol, 5-fluorouracil, methotrexate, dexamethasone, hydrocortisone, hydrocortisone acetate, hydroxy hydrocortisone, hydrocortisone valerate, triamcinolone acetonide, triamcinolone hexacetonide, amcinonide, flucinolone acetonide, flucinolone, flurandrenolide, difluorason diacetate, betamethasone dipropionate, betamethasone, betamethasone benzoate, betamethasone valerate, halcinonide, desoximethasone, desonide, prednisolone, clo cortolone pivalate, haloprogyn, iodochloro, miconazole nitrate, tolnaftate, thiabendazole, chloroxine, amphotericin, candicin, fungimycin, nystatin, chlordanol, clotrimazole, ethonam nitrate, miconazole nitrate, pyrrolnitrin, lezatione, ticlatone, tolnaftate, triacetin, dithiocarbamate, thiourea, thiocyanates; aromatic carboxylic acids and the amides thereof, benzoic acid, salicylic acid, salicylic acid amide and anilide; aromatic sulfides, polysulfides, and sulfoxides, 5,5-dichloro-2,2-dihydroxydiphenylsulfide; quaternary ammonia and phosphonium compounds, decamethylene-bis-(4-thiopyridine-methyl-tosylate); 8-hydroxyquinoline sulfate, halogenated quinolines, 7-iodo-8-hydroxyquinoline-5-sulfonic acid, 5-chloro-7-iodo-8-hydroxyquinoline, 5-chloro-8-hydroxyquinoline, 5,7-dichloro-8-hydroxyquinoline, 5,7-diiodo-8-hydroxyquinoline, decamethylene-bis-(4-aminoquinaldium chloride); (2-dimethylamino-6-(beta-diaminoethoxy)-benzothiazole dihydrochloride; 1-(o-chloro-alpha-

alpha-diphenyl-benzyl)-imidazole, 1-[o,p-dichloro-beta-(o,p-dichlorobenzoyloxy)-phenethylimidazole]; 2-phenylbenzimidazole, 2-furfurylbenzimidazole; 3,5-dibenzyltetrahydro-1,3,5-thiadiazine-2-thione; 5-nitro-2-furfuryl-3-chloropropionate; quinones, tetrachloro-p-benzoquinone, 1,4-naphthoquinone, phenanthraquinone; sulfonamide sulfones; aromatic diamines, 2-hydroxystilbarmidine, diamidinodiphenylamine, dioctyl sodium sulfosuccinate, miconazole nitrate, potassium sorbate, calcium propionate and sodium propionate, sodium lauryl sulfate, penicillin, cephalosporin, penicillinase, lincomycins, vancomycin, tetracycline, chloramphenicol, streptomycin, gramicidin, neomycin, polymyxin beta sulfate, tetracycline, benzethonium chloride, gentamicin sulfate, nitrofurazone, benzalkonium chloride, hexylresorcinol, chloroxylenol, cloflucaban, carboic acid (phenol), triclocaban, triclosan, glacial acetic acid, ascorbic acid, calcium pantothenate, lactic acid, salicylic acid, cantharidin, podophyllin, acyclovir, benzalkonium chloride, alcohol, allantoin, anhydrous glycerin, benzocaine, camphor, carbamide peroxide, lanolin, menthol, petrolatum, phenol, idoxuridine, amantadine, methasazone, cytarabine, interferon, chloroform, ether, bacillus calmette-guerin, levamisole, benzimidazole, tetramisole, levamisole, oxyquinoline, lidane, pyrethrin, piperonyl butoxide, malathion, crotamiton, benzoyl peroxide, resorcinol monoacetate, sulfur, povidone-iodine, phenol, fluocinolone acetonide, para-aminobenzoic acid, sodium thiosulfate, meclocycline sulfosalicylate, sodium sulfacetamide, tetracycline hydrochloride, 6-2 carbon aliphatic dicarboxylic acids, sulfated lime, coal tar, juniper tar, pine tar, allantoin, saponated cresol, menthol, mercury oleate, phenol, methotrexate 4-4'-diaminodiphenyl sulfone, benzocaine, lidocaine, lidocaine hydrochloride, dibucaine, dibucaine hydrochloride, procaine, tetracaine, tetracaine hydrochloride, tronothane, dyclonine hydrochloride, pramoxine hydrochloride, benzyl alcohol, diperodon, bulamben picrate, cyclomethycaine sulfate, dimethisoquin hydrochloride, N,N-dimethyl aspartic acid; N,N-dimethyl glutamic acid, trolamine salicylate, methyl salicylate; antipyrine, salicylamide, camphor, eugenol, eucalyptol, thymol, allyl isothiocyanate (mustard oil), capsicum preparations, histamine dihydrochloride, methyl nicotinate, turpentine oil, diphenhydramine hydrochloride, phenyltoloxamine dihydrochloride, pyrilamine maleate, triphenylamine hydrochloride, tetanus antigen, diphtheria antigen, streptococcus antigen, tuberculin, Candida antigen, Trichophyton antigen, Proteus antigen, vitamins B₃, B₅, B₆, A, D, and E, elastin, pregnenolone, hydroquinone, monobenzene, dioxybenzene, oxybenzene, padimate O, padimate A, aminobenzoic acid, cinoxate, diethanolamine p-methoxycinnamate, ethyl 4-bis(hydroxypropyl) aminobenzoate, ethylhexyl salicylate, glyceryl aminobenzoate, homosalate, lawson, dihydroxyacetone, red petrolatum, and sulisobenzonate.

The foregoing examples are illustrative of the methods and compositions of this invention and are not intended to be limiting. The present invention is believed to be a pioneering invention in the area of drug potentiation, and as such embraces many equivalents not specifically described herein.

Claims

1. A topical pharmaceutical composition comprising a pharmacologically active agent and a vehicle characterized in that said composition includes a potentiator for the pharmacologically active agent which comprises a water soluble zinc-containing compound that provides zinc ions that can complex or chelate with the pharmacologically active agent.
2. The pharmaceutical composition of claim 1, wherein the vehicle includes a carrier.
3. The pharmaceutical composition of claim 2, wherein the carrier is at least one member selected from the group consisting of water, acetone, isopropyl alcohol, stearyl alcohol, freons, ethyl alcohol, polyvinyl pyrrolidone, propylene glycol, polyethylene glycol, fragrance, gel-producing materials, mineral oil, stearic acid, spermactol, sorbitan monooleate, polysorbate, TWEEN, sorbitol, and methyl cellulose.
4. The pharmaceutical composition of claim 1, wherein the pharmacologically active agent and potentiator are provided in equimolar amounts and the composition is substantially free of other compounds that chelate or complex zinc.
5. The pharmaceutical composition of claim 1, wherein the potentiator is provided in an amount effective to chelate or complex the pharmacologically active agent.
6. The pharmaceutical composition of any preceding claim, further comprising at least one additive selected from the group consisting of colouring agents, stability-enhancing agents and antioxidants.
7. The pharmaceutical composition of any preceding claim, prepared in a dosage form.
8. The pharmaceutical composition of claim 7, prepared in a dosage form selected from the group consisting of lotion, ointment, cream, gel, suppository, nasal solution, mouthwash, spray, and aerosol.

9. The pharmaceutical composition of claim 1, wherein the potentiator is selected from the group consisting of zinc chloride, zinc iodide, zinc bromide, zinc sulfate, zinc nitrate, zinc stearate, and zinc acetate.
10. The pharmaceutical composition of any preceding claim, wherein the potentiator is zinc chloride.
11. The pharmaceutical composition of any preceding claim, wherein the pharmacologically active agent is selected from the group consisting of an antineoplastic agent, immunopharmacological agent, steroid, antiviral agent, anti-inflammatory agent, antifungal agent, antiparasitic agent, pediculicide, acne treatment compound, antiporiasis agent, leprosy agent, anaesthetic agent, counterirritant, antihistamine, allergy diagnostic agent, vitamin, nutrient, cosmetic agent, sunscreen, anti-unicellular microorganism agent, antimetabolite, cell-regulatory agent, allergen, analgesic agent, radiopaque agent, cryoprotective agent, perfume, insect repellent, hair dye, anticarring agent, melanin-stimulating agent, antiperspirant, antisecretory agent, depilatories, hair restorers, wrinkle reducing agent, anticandruft agent, emollients, rubifacient, and antibacterial agent.
12. The pharmaceutical composition of any preceding claim, wherein the pharmacologically active agent is steroid.
13. The pharmaceutical composition of any of claims 1 to 10, wherein the pharmacologically active agent is an antifungal agent.
14. The pharmacological composition of any of claims 1 to 10, wherein the pharmacologically active agent is an antibacterial agent.
15. The pharmacological composition of any of claims 1 to 10, wherein the pharmacologically active agent is selected from the group consisting of VP-16 (epidodophyllotoxin beta-D ethylidene glucopyranoside-etoposide); VM-26 (epidodophyllotoxin beta-D ethylidene glucopyranoside-teniposide); 4'-dimethyl epidodophyllotoxin; diethylstilbestrol; dithranol; cyclophosphamide; mitomycin; daunomycin; platinum cis-diamine-dichloride; adriamycin; altopurinol, 5-fluorouracil, methotrexate, and NDGA (nordihydroguaiaretic acid).
16. The pharmaceutical composition of any of claims 1 to 10, wherein the pharmacologically active agent is NDGA (nordihydroguaiaretic acid).
17. The pharmaceutical composition of any of claims 1 to 10, wherein the pharmacologically active agent is hydrocortisone or betamethasone.
18. The pharmaceutical composition of any of claims 1 to 10, wherein the pharmacologically active agent is vitamin A and its analogues or vitamin D and its analogues.
19. The pharmaceutical composition of any of claims 1 to 10, wherein the pharmacologically active agent is lincomycin.
20. The pharmaceutical composition of any of claims 1 to 10, wherein the pharmacological active agent is selected from the group consisting of polypeptide nanoparticles comprising interleukin or active fragments thereof, antibodies or active fragments thereof, interferons, liposomes; dexamethasone; hydrocortisone acetate; hydroxy hydrocortisone; hydrocortisone valerate; triamcinolone acetate; triamcinolone hexacetate; amcinonide; fluocinolone acetate; fluocinonide; flurandrenolide, difluorason diacetate; betamethasone dipropionate; betamethasone; betamethasone benzoate; betamethasone valerate; halcinonide; desoximethasone; desonide; prednisolone; dexamethasone pivalate; haloprogin, iodochloro, miconazole, nitrate, tolnafate, thiabendazole, chloroxine, amphotericin, candin, fungimycin, nystatin, chlordanol, clotrimazole, ethonam nitrate, miconazole nitrate, pyrrolnitrin, fezatione, ticlatone, tolnafate, triacetin, carbonic acid derivatives; dithiocarbamate, thiourea, thiocyanates; aromatic carboxylic acids and the amides thereof, benzoic acid, salicylic acid, salicylic acid amide and anilide; aromatic sulfides, polysulfides, and sulfoxides, 5,5-dichloro-2,2-dihydroxydiphenylsulfide; invert soaps, quaternary ammonia and phosphonium compounds, decamethylene-bis-(4-thio-pyridine-methyl-tosylate); quinoline derivatives, 8-hydroxy-quinoline sulfate, halogenated quinolines, 7-iodo-8-hydroxy-quinoline-5-sulfonic acid, 5-chloro-7-iodo-8-hydroxy-quinoline, 5-chloro-9-hydroxy-quinoline, 5,7-dichloro-8-hydroxyquinoline, 5,7-dichloro-8-hydroxyquinoline, decamethylene-bis (4-amino-quinidium chloride); benzothiazole derivatives, (2-dimethylamino-6-(beta-diaminoethoxy)-benzothiazole dihydrochloride); imidazole derivatives, 1-(o-chloro-alpha-phenylphenyl-benzyl)-imidazole, 1-[o,p-dichloro-beta-(o,p-dichlorobenzoyloxy)-phenethyl]-imidazole; benzimidazole derivatives, 2-phenylbenzimidazole, 2-furfurylbenzimidazole; thiazidine derivatives, 3,5-dibenzyltetrahydro-1,3,5-thiadiazine-2-thione; furan derivatives, 5-nitro-2-furfuryl-3-chloropropionate; quinones, tetrachloro-p-benzoquinone, 1,4-naphthoquinone; phenanthraquin-

none; sulfonamides and sulfones; aromatic diamidines, 2-hydroxystilbamidine, diaminodiphenylamine, dioctyl sodium sulfosuccinate, haloprogin, miconazole nitrate, potassium sorbate, propionate compounds, such as calcium propionate and sodium propionate, sodium lauryl sulfate; sulfonamides, penicillins, cephalosporins, penicillinase, lincomycins, vancomycins, tetracyclines, chloramphenicols, and streptomycins; gramicidin; neomycin; polymyxin beta sulfate; tetracycline; benzethonium chloride; gentamicin sulfate; nitrofurazone; benzalkonium chloride; hexylresorcinol; chloroxylenol; cloflucarban; carbolic acid (phenol); tridocarbon; triclosan; glacial acetic acid; ascorbic acid; calcium pantothenate; lactic acid; salicylic acid; cantharidin; podophyllin; acyclovir; benzalkonium chloride; alcohol, allantoin; anhydrous glycerin; benzocaine; camphor; carbamide peroxide; lanolin; menthol; petrolatum; phenol; urea; idoxuridine; amantadine; methisazone; cytarabine; interferone; chloroform; ether; bacillus calmette-guerin; levamisole; antiparasitic agent, anthelmintic agent, benzimidazole compounds; tetramisole; levamisole; isouquinoline compounds; lidane; pyrethrins; piperonyl butoxide; malathion; crotonitron; benzoyl peroxide; resorcinol; resorcinol monoacetate; sulfur; povidone-iodine; salicylic acid; phenol; fluocinolone acetonide; para-amino benzoic acid; sodium thiosulfate, mecloxyline sulfosalicylate; sodium sulfacetamide; tetracycline hydrochloride; aliphatic dicarboxylic acids, sulfated lime, cytosolic agent; keratolytic agent; tar preparation; hydrocortisone preparation; anti-itch preparation; antimicrobial; coal tar preparation; juniper tar preparation, pine tar preparation; allantoin; saponated cresol; menthol; mercury oleate; phenol preparation; resorcinol; salicylic acid; anthralin; methotrexate; 4,4'-diaminodiphenylsulfone; benzocaine; lidocaine; lidocaine hydrochloride; dibucaine; dibucaine hydrochloride; procaine; tetracaine; tetracaine hydrochloride; tronothane; dyclonine; dyclonine sulfate; dimethisoquin hydrochloride; salicylic acid derivatives; N,N-dimethyl aspartic acid; N,N-dimethyl glutamic acid; trolamine salicylate; methyl salicylate; antipyrine; aspirin; salicylamide; methyl salicylate; camphor; menthol; eugenol; eucalyptol; thymol; allyl isothiocyanate (mustard oil); capsicum preparations; histamine dihydrochloride; methyl nicotinate; turpentine oil; diphenhydramine hydrochloride; phenyltoloxamine dihydrogen citrate; pyrilamine maleate; triphenylamine hydrochloride; vitamin B₃, vitamin B₅, vitamin E; a vitamin or nutrient for skin, hair or scalp conditions; an anti-scaling agent; a cosmetic agent; a perfume; a composition to reduce the appearance of wrinkles; water soluble elastin; pregnenolone, skin depigmenting agents; bleaches; hydroquinone; monobenzone; dioxybenzone; oxybenzone; padimate O; padimate A; aminobenzoic acid; cinoxate; diethanolamine p-methoxycinnamate; ethyl 4-[bis (hydroxypropyl)] aminobenzoate; ethylhexyl; salicylate; glyceryl aminobenzoate; homosalate; the combination of lawsone and dihydroxyacetone; red petrolatum; sulisobenzene; allergenic extracts for diagnosis and immunotherapy of specific allergy offenders from the following categories; pollens, foods, dusts, epidermals, insects and stinging insects, fungi, molds, yeasts; tests for sensitivity to therapeutic penicillin (benzyl-penicilloyl-polylysine); tests for sensitivity to tetanus antigens, diptheria antigens; streptococcus antigens; tuberculin, Candida antigens, Trichophyton antigens, and Proteus antigens.

21. A topical pharmaceutical composition as in any of the preceding claims comprising a pharmacologically active agent and a vehicle characterized by a potentiator for the pharmacologically active agent which comprises a water soluble zinc-containing compound that provides zinc ions that can complex or chelate with the pharmacologically active agent for use in a treatment of dermatological disorders.

Skin Distribution of
(14)C-Nordihydroguaiaretic acid (NDGA) in Rats
Receiving (14)C-NDGA with and without ZnCl_2

